

CALIFORNIA STATE SCIENCE FAIR 2015 PROJECT SUMMARY

Name(s) **Project Number Elizabeth Fletes; Alex Herrera; Armeen Mobasher** S1709 **Project Title Observational Study of Properties of Active Galactic Nuclei** Abstract **Objectives/Goals** The objective of this project was to see how the morphology of galaxies who host Active Galactic Nuclei (AGN) are different from those that do not. Methods/Materials -If the Hardness Ratio of the sample is greater than -0.3 it is an AGN -If the Hardness Ratio of the sample is less than -0.3 it is a non-AGN -Gather additional data required to run the tests. Required data consists of: -RA, Dec, Full flux, HR Classical, Flux RADIUS, MAG AUTO, A IMAGE, B IMAGE Inputting Data into Galfit Using Aquamacs Input data into corresponding location in the Image and Galfit Control Parameters. Change path A to the location of the data image. (This is what you will be experimenting with) Change path B to the location of the output image. (This is where the results will end up in) Change path C to the location of the data image. Keep path D the same. Change path H to the size of the image. Change path I to the size of the convolution box (x, y). Change J to the magnitude (MAG AUTO). Sersic NOTE: Any unmentioned steps should remain unchanged Change #1 to the position of the AGN/Non-AGN. Change #2 to a number between 20-25. This number is about trial and error- there is no way to know exactly which number to input. Change #3 to Flux RADIUS. Keep #4 the same. Change #5 to a number between 0-4. Note: cannot equal 0. Subtract 90 from the position angle and input it into step #10. Sky In section sky, only #1 needs to be changed. To do so open the file in Topcat. When in the table, look for the header called "BACKGROUND" then type the corresponding number into #1. Running the Program with Galfit Open Terminal and type in "galfit" to open up galfit. Enter the name of the program to initiate the testing process. Results It was found that most of the galaxies were elliptical galaxies because their sersic_index was over 2.5. Based on the distribution of sersic parameters, it was found that there is a high likelihood for black holes to be hosted at the core of elliptical galaxies and not spirals. **Conclusions/Discussion Summary Statement** We were curious to see if the presence of a Black Hole at the center of a galaxy affected the shape and size of the galaxy. **Help Received** Under supervision of Dr. Mobasher, we were able to do our project at University of California Riverside. Our mentors who guided us were Laura Green and Vivian U.